



SELECTED COLUMNAR SECTIONS FOR THE COASTAL PLAIN, APPALACHIAN PLATEAUS, INTERIOR LOW PLATEAUS, AND VALLEY AND RIDGE PROVINCES IN ALABAMA

LAMAR COUNTY

SYSTEM	SERIES	GROUP, FORMATION, AND MEMBER		LITHO- LOGY	THICKNESS (feet)		THICKNESS (meters)	
Cretaceous	Upper	Tuscaloosa Group	Gordo Formation		449	98	137	30
			Coker Formation			351		107
Pennsylvanian	Lower	Pottsville Formation			2255		688	
Mississippian		Parkwood Formation			545		166	
		Bangor Limestone and Floyd Shale undifferentiated			393		120	
		Tuscumbia Limestone and Fort Payne Chert undifferentiated			121		37	
Devonian		Unnamed Devonian carbonate and chert unit			5		2	
Silurian		Silurian undifferentiated			645		197	
Ordovician		Ordovician undifferentiated			246		75	
		Ordovician undifferentiated			1130P		344P	

GEOLOGICAL SURVEY OF ALABAMA

Ernest A. Mancini
State Geologist

STRATIGRAPHY AND PALEONTOLOGY DIVISION


Charles W. Copeland
Chief Geologist

ATLAS SERIES 20

**SELECTED COLUMNAR SECTIONS FOR THE COASTAL PLAIN,
APPALACHIAN PLATEAUS, INTERIOR LOW PLATEAUS, AND
VALLEY AND RIDGE PROVINCES IN ALABAMA**

By

Dorothy E. Raymond and Charles W. Copeland



GEOLOGICAL SURVEY OF ALABAMA

420 Hackberry Lane
P. O. Box 0
Tuscaloosa, Alabama 35486 - 9780
(205) 349-2852

DIRECTORS
Executive Assistant, R. C. Bagenstose
Stratigraphy &
Paleontology, C. W. Copeland, Jr.
Mineral Resources, W. E. Smith
Energy Resources, M. L. Epsman
Water Resources, J. D. Moore
Biological Resources, M. F. Mettee
Geochemistry Laboratory, N. A. Lloyd
Geocartography, T. V. Stone

October 23, 1987

Honorable Guy Hunt
Governor of Alabama
Montgomery, Alabama

Dear Governor Hunt:

I have the honor to transmit herewith a report entitled "Selected Columnar Sections for the Coastal Plain, Appalachian Plateaus, Interior Low Plateaus, and Valley and Ridge Provinces in Alabama" by Dorothy E. Raymond and Charles W. Copeland, which has been printed as Atlas Series 20 of the Geological Survey of Alabama.

The report graphically illustrates the stratigraphy of the state through columnar sections showing rock units and assigned formations encountered by oil, gas and water wells that have been drilled in Alabama. The details of lithology, age, and thickness of individual formations should make this publication useful in the exploration for water, mineral and energy resources.

The columnar sections selected for inclusion in the atlas typify the formations occurring in the different parts of the state and the facies changes that occur. Also included are columnar sections of some of the newer wells that have been drilled in previously unexplored areas.

Respectfully,

Ernest A. Mancini

Ernest A. Mancini
State Geologist

CONTENTS

	Page
Introduction	1
Acknowledgments	1
References	1
Index maps	2
Legend	3
Coastal Plain province	5
Colbert County (Column 1)	7
Franklin County (Column 2)	7
Lamar County (Column 3)	9
Pickens County (Column 4)	9
Greene County (Column 5)	11
Sumter County (Column 6)	11
Choctaw County (Column 7)	13
Washington County (Column 8)	15
Mobile County (Column 9)	17
Mobile Bay (Column 10)	19
Baldwin County (Column 11)	21
Covington County (Column 12)	23
Dallas County (Column 13)	23
Houston County (Column 14)	25
Barbour County (Column 15)	25
Appalachian Plateaus province and Interior Low Plateaus province	27
Madison County (Column 16)	29
Jackson County (Column 17)	29
De Kalb County and western Etowah County (Column 18)	31
Blount County (Column 19)	31
Cullman County (Column 20)	33
Morgan County (Column 21)	33
Valley and Ridge province	35
Jefferson County (Column 22)	37
Northeastern Etowah and northwestern Cherokee Counties (Column 23)	37
Calhoun, southeastern Cherokee, and southeastern Etowah Counties (Column 24)	39
Shelby County (Column 25)	41
Bibb and Shelby Counties (Column 26)	43

INTRODUCTION

This atlas is a compilation of stratigraphic columns for selected areas of Alabama and is designed to summarize the stratigraphy of the state, exclusive of the igneous and metamorphic rocks of the Piedmont physiographic province. Locations of columns were selected to illustrate facies changes and thickness changes across the state. Research for most of the columns was initiated during the preparation of stratigraphic columns for the COSUNA (Correlation of Stratigraphic Units for North America) Project under the auspices of the American Association of Petroleum Geologists.

The columns are based on data from oil and gas test wells, water wells, published literature, and open-file reports. In some areas of the state considerable information is available either owing to better outcrops or more test holes; consequently, columns for these areas are more detailed. In areas where outcrops are few and scattered and where test drilling is limited, the columns are more generalized and represent larger areas (two or more counties).

Specific sources of information used to generate each column are given below the respective column. Selected references on the area of concern are given on the page facing the column.

ACKNOWLEDGMENTS

The authors wish to express appreciation to colleagues, Gary V. Wilson, Assistant Oil and Gas Supervisor, State Oil and Gas Board (Alabama), for helpful suggestions and assistance concerning formation tops in wildcat oil and gas test wells, and W. Edward Osborne, stratigrapher, Geological Survey of Alabama, for invaluable assistance with thickness data on formations in the Valley and Ridge and Plateaus.

REFERENCES


Copeland, C. W., 1985, Column 29, *in* Shaver, R. H., regional coordinator, Midwestern Basin and Arches region, Correlation of Stratigraphic Units of North America (COSUNA) Project: American Association of Petroleum Geologists Correlation Chart Series, 1984.

_____, 1987, Column 19, *in* Mankin, C. J., regional coordinator, Texas-Oklahoma tectonic region, Correlation of Stratigraphic Units of North America (COSUNA) Project: American Association of Petroleum Geologists Correlation Chart Series, 1986.

Copeland, C. W., and Raymond, D. E., 1985, Columns 1 through 8, *in* Patchen, D. G., Avary, K. L., and Erwin, R. B., regional coordinators, Southern Appalachian region, Correlation of Stratigraphic Units of North America (COSUNA) Project: American Association of Petroleum Geologists Correlation Chart Series, 1984.

_____, in press, Columns 15 through 20, *in* Gulf Coast region, Correlation of Stratigraphic Units of North America (COSUNA) Project: American Association of Petroleum Geologists Correlation Chart Series.

INDEX MAPS

































EXPLANATION

- 2 Column number
- Location of column


The locations represented by columns are indicated on the above index maps. Locations are shown by county and physiographic province. All columns are located in the Coastal Plain, Valley and Ridge, Appalachian Plateaus, and Interior Low Plateaus physiographic provinces, which are underlain by sedimentary rocks. No columns were attempted for the Piedmont province because well data in this area are generally lacking.

Although most columns are generalized to represent a large area, many columns are based in large part on stratigraphy in specific oil and gas test wells. In these cases, locations shown on the index map represent the locations of these specific wells. In some cases, however, columns are composited entirely on the basis of stratigraphy in scattered outcrops over a two or three county area. In these cases, primarily in the Valley and Ridge province, locations given on the index map represent hypothetical locations, generally near the centers of the areas of concern.

LEGEND

	Gravel or conglomerate		Silt, siltstone, or silty shale		Sandy dolomite
	Sand or sandstone		Shale		Cherty dolomite
	Argillaceous sandstone		Sandy shale		Anhydrite
	Calcareous sandstone		Calcareous shale		Anhydritic shale
	Limestone or chalk		Carbonaceous shale		Anhydritic limestone
	Argillaceous limestone		Interbedded sandstone and shale		Bentonite
	Sandy limestone		Interbedded shale and limestone		Salt
	Oolitic limestone		Dolomite		Coal or lignite
	Cherty limestone		Dolomitic limestone or calcareous dolomite		Basement rock
	Bedded chert		Argillaceous limestone		Thrust fault

E Estimated
P Penetrated
Exp Exposed



Vertical Scale

COASTAL PLAIN PROVINCE

SELECTED REFERENCES

Column 1	Column 2
Harris, H. B., Moore, G. K., and West, L. R., 1963, Geology and ground-water resources of Colbert County, Alabama: Alabama Geological Survey County Report 10, 71 p.	Jewell, J. W., 1969, An oil and gas evaluation of north Alabama: Alabama Geological Survey Bulletin 93, 65 p.
Henry, T. W., Gordon, MacKenzie, Jr., Schweinfurth, S. P., and Gillespie, W. H., 1985, Significance of the goniatite <i>Bilinguites eliasi</i> and associated biotas, Parkwood Formation and Bangor Limestone, northwestern Alabama: Journal of Paleontology, v. 59, no. 5, p. 1138-1145.	Kidd, J. T., 1975, Pre-Mississippian subsurface stratigraphy of the Warrior Basin in Alabama: Gulf Coast Association Geological Societies Transactions, v. 25, p. 20-39.
Jewell, J. W., 1969, An oil and gas evaluation of north Alabama: Alabama Geological Survey Bulletin 93, 65 p.	Kidd, J. T., and Neathery, T. L., 1976, Correlation between Cambrian rocks of the southern Appalachian geosyncline and Interior Low Plateaus: Geology, v. 4, p. 767-769.
Kidd, J. T., 1975, Pre-Mississippian subsurface stratigraphy of the Warrior Basin in Alabama: Gulf Coast Association Geological Societies Transactions, v. 25, p. 20-39.	Peace, R. R., Jr., 1963, Geology and groundwater resources of Franklin County, Alabama, a reconnaissance report: Alabama Geological Survey Bulletin 72, 55 p.
Kidd, J. T., and Neathery, T. L., 1976, Correlation between Cambrian rocks of the southern Appalachian geosyncline and Interior Low Plateaus: Geology, v. 4, p. 767-769.	Thomas, W. A., 1972, Mississippian stratigraphy of Alabama: Alabama Geological Survey Monograph 12, 121 p.
Thomas, W. A., 1972, Mississippian stratigraphy of Alabama: Alabama Geological Survey Monograph 12, 121 p.	

COLUMN 1
COLBERT COUNTY

SYSTEM	SERIES	GROUP, FORMATION AND MEMBER	LITHO-LOGY	THICKNESS (feet)	THICKNESS (meters)
Cretaceous	Upper	Tuscaloosa Group		50	15
		Bangor Limestone		90	27
Mississippian	Upper	Hartselle Sandstone		140	43
		Pride Mountain Formation		260	79
		Tuscumbia Limestone		200	61
		Fort Payne Chert		183	56
	Lower	Chattanooga Shale		22	7
Devonian		Unnamed Silurian and Devonian carbonate		180	55
Silurian		Wayne Group		83	25
Ordo- vician	Upper	Maysville and Nashville Groups undifferentiated		240	73
	Middle	Stones River Group		639	195
	Lower				
Cambrian	Upper	Knox Group undifferentiated		3500 E	1067 E
		Ketona Dolomite		1100 E	335 E
	Middle	Conasauga Formation		500 E	152 E
		Basement complex			

From Lower Ordovician to Upper Cretaceous, the Colbert County section is based on cited references, outcrops, and the sample description of an oil and gas test well (Alabama State Oil and Gas Board Permit Number 576). The Precambrian to Lower Ordovician section is based on data from wells drilled in Cullman and Morgan Counties, Alabama. Total depth of well Permit Number 576 is 1,978 feet.

COLUMN 2
FRANKLIN COUNTY

SYSTEM	SERIES	GROUP, FORMATION AND MEMBER	LITHO-LOGY	THICKNESS (feet)	THICKNESS (meters)
Cretaceous	Upper	Tuscaloosa Group		50	15
		Bangor Limestone		650	198
Pennsylvanian	Lower	Pottsville Formation		197	60
		Parkwood Formation		222	67
Mississippian	Upper	Hartselle Sandstone		57	17
		Pride Mountain Formation		283	86
		Tuscumbia Limestone		125	38
		Fort Payne Chert		125	38
	Lower	Chattanooga Shale		20	6
Devonian		Unnamed Devonian carbonate and chert unit		62	19
Silurian		Silurian undifferentiated		285	87
Ordo- vician	Upper	Maysville and Nashville Groups undifferentiated		233	71
	Middle	Stones River Group		540	165
	Lower				
Cambrian	Upper	Knox Group undifferentiated		3500 E	1067 E
		Ketona Dolomite		1100 E	335 E
	Middle	Conasauga Formation		750 E	229 E
		Basement complex			

From Lower Ordovician to Upper Cretaceous, the Franklin County section is based on cited references, outcrops and the sample description of an oil and gas test well (Alabama State Oil and Gas Board Permit Number 2051). The Precambrian-Lower Ordovician section is based on data from wells drilled in Cullman and Morgan Counties, Alabama. The total depth of well Permit Number 2051 is 4,000 feet.

SELECTED REFERENCES

Column 3

Drennen, C. W., 1953, Stratigraphy and structure of outcropping pre-Selma Coastal Plain beds of Fayette and Lamar Counties, Alabama: U.S. Geological Survey Circular 267, 9 p.

Kidd, J. T., 1975, Pre-Mississippian subsurface stratigraphy of the Warrior Basin in Alabama: Gulf Coast Association Geological Societies Transactions, v. 25, p. 20-39.

Smith, W. E., 1979, Pennsylvanian stratigraphy of Alabama, *in* The Mississippian and Pennsylvanian (Carboniferous) Systems in the United States: U.S. Geological Survey Professional Paper 1110 I, p. 123-136.

Thomas, W. A., 1972, Mississippian stratigraphy of Alabama: Alabama Geological Survey Monograph 12, 121 p.

Column 4

Kidd, J. T., 1975, Pre-Mississippian subsurface stratigraphy of the Warrior Basin in Alabama: Gulf Coast Association Geological Societies Transactions, v. 25, p. 20-39.

Smith, W. E., 1979, Pennsylvanian stratigraphy of Alabama, *in* The Mississippian and Pennsylvanian (Carboniferous) Systems in the United States: U.S. Geological Survey Professional Paper 1110 I, p. 123 - 136.

Thomas, W. A., 1972, Mississippian stratigraphy of Alabama: Alabama Geological Survey Monograph 12, 121 p.

Wahl, K. D., 1965, Ground-water resources of Pickens County, Alabama, a reconnaissance: Alabama Geological Survey Bulletin 83, 84 p.

COLUMN 3
LAMAR COUNTY

SYSTEM	SERIES	GROUP, FORMATION, AND MEMBER		LITHO-LOGY	THICKNESS (feet)		THICKNESS (meters)	
Cretaceous	Upper	Tuscaloosa Group	Gordo Formation		449	98	137	30
			Coker Formation			351		107
Pennsylvanian	Lower	Pottsville Formation			2255	688		
Mississippian		Parkwood Formation			545	166		
		Bangor Limestone and Floyd Shale undifferentiated			393	120		
		Tuscumbia Limestone and Fort Payne Chert undifferentiated			121	37		
Devonian		Chattanooga Shale			5	2		
		Unnamed Devonian carbonate and chert unit			645	197		
Silurian		Silurian undifferentiated			246	75		
Ordovician		Ordovician undifferentiated			1130P	344P		

The Lamar County section is based on cited references and data from an oil and gas test well (Alabama State Oil and Gas Board Permit Number 2527). Total depth of the well is 5,990 feet.

COLUMN 4
PICKENS COUNTY

SYSTEM	SERIES	GROUP, FORMATION, AND MEMBER		LITHO-LOGY	THICKNESS (feet)		THICKNESS (meters)	
Cretaceous	Upper	Eutaw Formation			785	270	82	
		Tuscaloosa Group	Gordo Formation			260	79	
			Coker Formation			525	160	
Pennsylvanian	Lower	Pottsville Formation			4350	1326		
Mississippian		Parkwood Formation			525	160		
		Bangor Limestone and Floyd Shale undifferentiated				422	129	
		Tuscumbia Limestone and Fort Payne Chert undifferentiated				77	23	
Devonian		Chattanooga Shale			14	4		
		Unnamed Devonian carbonate and chert unit				664	202	
Silurian		Silurian undifferentiated			225	69		
Ordovician		Ordovician undifferentiated			3208	978		
	Lower	Knox Group undifferentiated			449P	137P		
Cam- brian	Upper							

The Pickens County section is based on cited references and the sample description of an oil and gas test well (Alabama State Oil and Gas Board Permit Number 1634). Total depth of the well is 10,995 feet.

SELECTED REFERENCES

Column 5

Kidd, J. T., 1975, Pre-Mississippian subsurface stratigraphy of the Warrior Basin in Alabama: Gulf Coast Association Geological Societies Transactions, v. 25, p. 20-39.

Smith, W. E., 1979, Pennsylvanian stratigraphy of Alabama, *in* The Mississippian and Pennsylvanian (Carboniferous) Systems in the United States: U.S. Geological Survey Professional Paper 1110 I, p. 123 - 136.

Thomas, W. A., 1972, Mississippian stratigraphy of Alabama: Alabama Geological Survey Monograph 12, 121 p.

Wahl, K. D., 1966, Geology and ground-water resources of Greene County, Alabama: Alabama Geological Survey Bulletin 86, 93 p.


Column 6

Davis, M. E., Sanford, T. H., Jr., and Jefferson, P. O., 1980, Water availability and geology of Sumter County, Alabama: Alabama Geological Survey Special Map 158, 69 p.

Monroe, W. H., 1941, Notes on deposits of Selma and Ripley age in Alabama: Alabama Geological Survey Bulletin 48, 150 p.











Monroe, W. H., and Hunt, J. L., 1958, Geology of the Epes Quadrangle, Alabama: U.S. Geological Survey Geologic Quadrangle Map GQ-113.

Oxley, M. L., Minihan, Edward, and Ridgway, J. M., 1967, A study of the Jurassic sediments in portions of Mississippi and Alabama: Gulf Coast Association Geological Societies Transactions, v. 17, p. 24-48.

SYSTEM	SERIES	GROUP FORMATION, AND MEMBER		LITHO- LOGY	THICKNESS (feet)	THICKNESS (meters)
Cretaceous	Upper	Eutaw Formation			420	128
		Tuscaloosa Group	Gordo Formation		290	88
			Coker Formation		1095 805	333 245
	Lower	Lower Cretaceous undiff.			165	50
Pennsylvanian	Lower	Pottsville Formation		4835	1474	
		Parkwood Formation		665	203	
		Floyd Shale		1145	349	
Mississippian		Tuscumbia Limestone and Fort Payne Chert undifferentiated		115	35	
		Chattanooga Shale		80	24	
		Unnamed Devonian carbonate and chert unit		787	240	
Devonian		Silurian undifferentiated		248	76	
Ordovician		Ordovician undifferentiated		3090	942	
		Knox Group		210 +	64 +	
Cam- brian						

COLUMN 5
GREENE COUNTY

COLUMN 6
SUMTER COUNTY

SYSTEM	SERIES	GROUP, FORMATION, AND MEMBER	LITHO- LOGY	THICKNESS (feet)	THICKNESS (meters)
Cretaceous	Paleocene	Porters Creek Formation		246	75
		Clayton Formation		40	12
	Upper	Prairie Bluff Chalk		100	30
		Ripley Formation		175	53
		Demopolis Chalk		310	94
		Arcola Limestone Mbr.		10	3
		Mooreville Chalk		262	80
		Eutaw Formation		418	128
	Lower	Gordo Formation		445	136
		Coker Formation		795	242
		Lower Cretaceous undifferentiated		802	244
	Upper	Cotton Valley Group		438P	134P
	Lower				
Juras- sic	Upper				

The Sumter County section is based on the sample and electric logs of an oil and gas test well (Alabama State Oil and Gas Board Permit Number 445). Total depth of the well is 4,010 feet.

The Greene County section is based on the sample and electric logs of an oil and gas test well (Alabama State Oil and Gas Board Permit Number 1810). Total depth of the well is 12,850 feet.

SELECTED REFERENCES

Column 7

Moore, D. B., 1971, Subsurface geology of southwest Alabama: Alabama Geological Survey Bulletin 99, 80 p.

Tolson, J. S., Copeland, C. W., and Bearden, B. L., 1983, Stratigraphic profiles of Jurassic strata in the western part of the Alabama Coastal Plain: Alabama Geological Survey Bulletin 122, 425 p.

Turner, J. D., and Newton, J. G., 1971, Geology of Choctaw County, Alabama: Alabama Geological Survey Special Map 102, with text, 15 p.

COLUMN 7
CHOCTAW COUNTY

SYSTEM	SERIES	GROUP, FORMATION AND MEMBER	LITHO-LOGY	THICKNESS (feet)	THICKNESS (meters)
Tertiary	Eocene	Lisbon Formation and Gosport Sand undifferentiated		250	76
		Tallahatta Formation		180	55
		Wilcox Group undifferentiated		1352	412
	Paleocene	Naheola Formation		423	129
		Porters Creek Formation		362	110
		Clayton Formation		23	7
	Upper	Selma Group		716	117
		Eutaw Formation		384	218
		Tuscaloosa Group	"Upper Tuscaloosa"	600	183
			"Marine Tuscaloosa"	440	134
			"Lower Tuscaloosa"	563	172
				1603	489
				440	134
				563	172
Cretaceous	Lower	Lower Cretaceous undifferentiated		4064	1239











SYSTEM	SERIES	GROUP, FORMATION AND MEMBER	LITHO-LOGY	THICKNESS (feet)	THICKNESS (meters)
Cretaceous	Lower	Cotton Valley Group		2236	682
Jurassic	Upper	Haynesville Formation		939	286
		Buckner Anhydrite Member		225	69
		Smackover Formation		474	144
		Norphlet Formation		262	80
		Louann Salt		868	265
	Lower	Werner Formation		1362	415
Triassic		Eagle Mills Formation		1349 P	411 P




The Choctaw County column is based on outcrop and subsurface data. The Tertiary and uppermost Cretaceous is based primarily on the electric log and sample log from an oil and gas test well (Alabama State Oil and Gas Board Permit Number 502). The Cretaceous through Triassic section is based on the electric logs and sample log from an oil and gas test well (Alabama State Oil and Gas Board Permit Number 3589). Total depths of the two wells are 4,116 and 17,403 feet, respectively.

SELECTED REFERENCES

Column 8

- Moore, D. B., 1971, Subsurface geology of southwest Alabama: Alabama Geological Survey Bulletin 99, 80 p.
- Tolson, J. S., Copeland, C. W., and Bearden, B. L., 1983, Stratigraphic profiles of Jurassic strata in the western part of the Alabama Coastal Plain: Alabama Geological Survey Bulletin 122, 425 p.
- Turner, J. D., and Newton, J. G., 1971, Geologic map of Washington County, Alabama: Alabama Geological Survey Special Map 100, with text, 9 p.

SYSTEM	SERIES	GROUP, FORMATION, AND MEMBER		LITHO-LOGY	THICKNESS (feet)	THICKNESS (meters)
Tertiary	Miocene	Miocene undifferentiated			328	100
	Oligocene	Oligocene undifferentiated			216	66
	Eocene	Jackson Group			272	83
		Claiborne Group			728	222
	Paleocene	Wilcox Group			1312	400
		Naheola Formation			572	174
		Porters Creek Formation			473	144
		Clayton Formation			7	2
Cretaceous	Upper	Selma Group			1165	355
		Eutaw Formation			538	164
		Tuscaloosa Group	"Upper Tuscaloosa"		683	208
			"Marine Tuscaloosa"		197	60
	"Lower Tuscaloosa"			594	181	
	Lower	Lower Cretaceous undifferentiated			4918	1499

SYSTEM	SERIES	GROUP, FORMATION, AND MEMBER	LITHO-LOGY	THICKNESS (feet)	THICKNESS (meters)
Cretaceous	Lower	Cotton Valley Group		2466	752
Jurassic	Upper		Haynesville Formation		1686
		Buckner Anhydrite Member		151	46
		Smackover Formation		525	160
		Norphlet Formation		161P	49P

The Washington County section is a composite from adjacent oil and gas test wells. The Mesozoic section is from data interpreted from Alabama State Oil and Gas Board Permit Number 1997, and the Cenozoic section is based on the sample description of Alabama State Oil and Gas Board Permit Number 982. Total depths of the two wells are 16,950 and 12,209 feet, respectively.

SELECTED REFERENCES

Column 9

- Eaves, Everett, 1976, Citronelle Oil Field, Mobile County, Alabama: American Association Petroleum Geologists Memoir 24, p. 259-275.
- Maher, J. C., and Applin, E. R., 1968, Correlation of subsurface Mesozoic and Cenozoic rocks along the Eastern Gulf Coast: American Association Petroleum Geologists Cross Section Publication 6, p. 11 and 12.
- Mancini, E. A., and Payton, J. W., 1981, Petroleum geology of South Carlton Field, Lower Tuscaloosa "Pilot sand," Clarke and Baldwin Counties, Alabama: Gulf Coast Association Geological Societies Transactions, v. 31, p. 139-147.
- Moore, D. B., 1971, Subsurface geology of southwest Alabama: Alabama Geological Survey Bulletin 99, 80 p.
- Raymond, D. E., 1985, Depositional sequences in the Pensacola Clay (Miocene) of southwest Alabama: Alabama Geological Survey Bulletin 114, 87 p.
- Reed, P. C., 1971, Geologic map of Mobile County, Alabama: Alabama Geological Survey Special Map 93, with text, 22 p.
- Tolson, J. S., Copeland, C. W., and Bearden, B. L., 1983, Stratigraphic profiles of Jurassic strata in the western part of the Alabama Coastal Plain: Alabama Geological Survey Bulletin 122, 425 p.

COLUMN 9

MOBILE COUNTY

SYSTEM	SERIES	GROUP, FORMATION, AND MEMBER		LITHO-LOGY	THICKNESS (feet)	THICKNESS (meters)
Quaternary	Holocene	Coastal and low-level fluvial deposits			10 - 150	3 - 46
	Pleistocene	High-level fluvial deposits			13 - 40	4 - 12
		Citronelle Formation			0 - 197	0 - 60
Tertiary	Pliocene					
	Miocene	Miocene undifferentiated			1687	514
	Oligocene	Tampa Formation and Chickasawhay Limestone undifferentiated			193	59
		Bucatunna Clay Mbr. of Byram Formation			140	43
					170	52
	Eocene	Jackson Group				
		Claiborne Group	Lisbon Formation		160	49
			Tallahatta Formation		350	107
		Bashi Marl			10	3
	Paleocene	Wilcox Group			1750	533
		Salt Mountain Limestone			36	11
		Midway Group			1010	308
Cretaceous	Upper	Selma Group			1260	384
		Eutaw Formation			165	50
		Tuscaloosa Group	"Upper Tuscaloosa"		410	125
			"Marine Tuscaloosa"		362	110
			"Lower Tuscaloosa"		703	214
		Washita and Fredericksburg Groups undifferentiated				
	Lower				2106	642

SYSTEM	SERIES	GROUP, FORMATION, AND MEMBER		LITHO-LOGY	THICKNESS (feet)	THICKNESS (meters)
Cretaceous	Lower	Paluxy Formation			1246	380
		Mooringsport Formation			235	72
		Ferry Lake Anhydrite			233	71
		Rodessa Formation			417	127
		Sligo Formation			531	162
		Hosston Formation			1652	504
Jurassic	Upper	Cotton Valley Group			2431	741
		Haynesville Formation			1142	348
		Buckner Anhydrite Member			249	76
		Smackover Formation			44	13
		Norphlet Formation			406	124
	Middle	Louann Salt			542	165
		Werner Formation			92	28
Devonian	Upper	Basement complex (phyllite)			226 p	69 p

The information on Pliocene and younger units is from surface exposures and cited references. The remainder of the section is from the sample descriptions and other records of oil and gas test wells (Alabama State Oil and Gas Board Permit Numbers 1862 and 971. Thickness data on the Jurassic section are from Permit Number 1862. This well is near the crest of the Wiggins arch and has a total depth of 20,089 feet.

SELECTED REFERENCES

Column 10

Eaves, Everett, 1976, Citronelle Oil Field, Mobile County, Alabama: American Association Petroleum Geologists Memoir 24, p. 259-275.

Maher, J. C., and Applin, E. R., 1968, Correlation of subsurface Mesozoic and Cenozoic rocks along the Eastern Gulf Coast: American Association Petroleum Geologists Cross Section Publication 6, p. 11 and 12.

Mancini, E. A., Mink, R. M., Bearden, B. L., and Wilkerson, R. P., 1985, Norphlet Formation (Upper Jurassic) of southwestern and offshore Alabama: Environments of deposition and petroleum geology: American Association Petroleum Geologists Bulletin 69, no. 6, p. 881-898.

Mancini, E. A., and Payton, J. W., 1981, Petroleum geology of South Carlton Field, Lower Tuscaloosa "Pilot sand," Clarke and Baldwin Counties, Alabama: Gulf Coast Association Geological Societies Transactions, v. 31, p. 139-147.

Mink, R. M., Bearden, B. L., and Mancini, E. A., 1985, Regional Jurassic geologic framework of Alabama coastal waters area and adjacent Federal waters area: Alabama State Oil and Gas Board Report, no. 12, 58 p.

Moore, D. B., 1971, Subsurface geology of southwest Alabama: Alabama Geological Survey Bulletin 99, 80 p.

Pittman, J. G., 1985, Correlation of beds within the Ferry Lake Anhydrite of the Gulf Coastal Plain: Gulf Coast Association Geological Societies Transactions, v. 35, p. 251-260.

Raymond, D. E., 1985, Depositional sequences in the Pensacola Clay (Miocene) of southwest Alabama: Alabama Geological Survey Bulletin 114, 87 p.

Tolson, J. S., Copeland, C. W., and Bearden, B. L., 1983, Stratigraphic profiles of Jurassic strata in the western part of the Alabama Coastal Plain: Alabama Geological Survey Bulletin 122, 425 p.

COLUMN 10

MOBILE BAY

SYSTEM	SERIES	GROUP, FORMATION AND MEMBER	LITHO-LOGY	THICKNESS (feet)	THICKNESS (meters)
Tertiary	Holocene	Coastal deposits		40	12
	Pliocene				
	Miocene	Miocene undifferentiated		1086	331
		Pensacola Clay		1630	497
		Tampa Formation and Chickasawhay Limestone undifferentiated		432	132
		Bucatunna Clay Mbr. of Byram Formation		274	84
	Oligocene				
	Eocene	Jackson Group		50	15
		Claiborne Group	Lisbon Fm. Tallahatta Formation	268 76 192 81 23 58	
		Wilcox Group		1076	328
	Paleocene	Midway Group		942	287
Cretaceous	Upper	Selma Group		1402	427
		Eutaw Formation		148	45
		Tuscaloosa Group	"Upper Tuscaloosa"	408	124
			"Marine Tuscaloosa"	330	101
			"Lower Tuscaloosa"	788	240
		Washita and Fredericksburg Groups undifferentiated		1526	465
	Lower			2529	771

SYSTEM	SERIES	GROUP, FORMATION, AND MEMBER	LITHO-LOGY	THICKNESS (feet)	THICKNESS (meters)
Cretaceous	Lower	Paluxy Formation		725	221
		Mooringsport Formation		864	263
		Ferry Lake Anhydrite		293	89
		Rodessa Formation		669	204
		Sligo Formation		256	78
		Hosston Formation		2149	655
		Cotton Valley Group		2709	826
		Haynesville Formation		785	239
		Smackover Formation		711	217
		Norphlet Formation		622	190
Jurassic	Upper	"Pine Hill anhydrite" Louann Salt		37P	11P

The column is based on sample descriptions and electric logs for an oil and gas test well (Alabama State Oil and Gas Board Permit Number 2543). Interpretation of lithologies and stratigraphic units in the Lower Cretaceous was aided by the use of unpublished descriptions of a nearby well (Alabama State Oil and Gas Board Permit Number 3127) by H. V. Wuerch. Total depth of well Permit Number 2543 is 21,113 feet.

SELECTED REFERENCES

Column 11

Eaves, Everett, 1976, Citronelle Oil Field, Mobile County, Alabama: American Association Petroleum Geologists Memoir 24, p. 259-275.

Maher, J. C., and Applin, E. R., 1968, Correlation of subsurface Mesozoic and Cenozoic rocks along the Eastern Gulf Coast: American Association Petroleum Geologists Cross Section Publication 6, p. 11 and 12.

Mancini, E. A., and Payton, J. W., 1981, Petroleum geology of South Carlton Field, Lower Tuscaloosa "Pilot sand," Clarke and Baldwin Counties, Alabama: Gulf Coast Association Geological Societies Transactions, v. 31, p. 139-147.

Moore, D. B., 1971, Subsurface geology of southwest Alabama: Alabama Geological Survey Bulletin 99, 80 p.

Raymond, D. E., 1985, Depositional sequences in the Pensacola Clay (Miocene) of southwest Alabama: Alabama Geological Survey Bulletin 114, 87 p.

Reed, P. C., 1971, Geologic map of Baldwin County, Alabama: Alabama Geological Survey Special Map 94, with text, 5 p.

Tolson, J. S., Copeland, C. W., and Bearden, B. L., 1983, Stratigraphic profiles of Jurassic strata in the western part of the Alabama Coastal Plain: Alabama Geological Survey Bulletin 122, 425 p.

COLUMN 11
BALDWIN COUNTY

SYSTEM	SERIES	GROUP, FORMATION AND MEMBER		LITHO- LOGY	THICKNESS (feet)	THICKNESS (meters)
Tertiary	Holocene	Coastal and low-level fluvial deposits			10 - 150	3 - 46
	Pleistocene	High-level fluvial deposits			13 - 49	4 - 15
	Pliocene	Citronelle Formation			0 - 130	0 - 40
	Miocene	Miocene undifferentiated			669	204
		Pensacola Clay	Upper member		321	98
			Escambia Sand Mbr.		56	17
			Lower member		334	102
		Tampa Formation and Chickasawhay Limestone undifferentiated			315	96
	Oligocene	Bucaturunna Clay Mbr. of Byram Formation			184	56
		Jackson Group			256	78
	Eocene	Lisbon Formation			216	66
		Tallahatta Formation			341	104
		Wilcox Group			1161	354
	Paleocene	Midway Group			879	268
Cretaceous	Upper	Selma Group			1547	472
		Eutaw Formation			254	77
		Tuscaloosa Group	"Upper Tuscaloosa"		190	58
			"Marine Tuscaloosa"		535	163
			"Lower Tuscaloosa"		611	186
	Lower	Lower Cretaceous undifferentiated			1336	407

SYSTEM	SERIES	GROUP, FORMATION AND MEMBER	LITHO- LOGY	THICKNESS (feet)	THICKNESS (meters)
Cretaceous	Lower	Lower Cretaceous undifferentiated		7310	2228
Jurassic	Upper	Cotton Valley Group		1925	587
		Haynesville Formation		1926	587
	Middle	Smackover Formation		500	152
		Norphlet Formation		105	32
		Louann Salt		52P	16P

The information on Pliocene and younger units is from surface exposures. The remainder of the section is from the sample description and other records of an oil and gas test well (Alabama State Oil and Gas Board Permit Number 2339). Total depth of the well is 19,402 feet.

SELECTED REFERENCES

Column 12

Huddlestun, P. F., 1965, Correlation of upper Eocene and lower Oligocene strata between the Sepulga, Conecuh, and Choctawhatchee Rivers of southern Alabama: Florida State University Master's thesis, 101 p.
Turner, J. D., and Scott, J. C., 1968, Geologic map of Covington County, Alabama: Alabama Geological Survey Special Map 66.

Column 13

Scott, J. C., Golden, H. G., and Newton, J. G., 1981, Geology and water availability of Dallas County, Alabama: Alabama Geological Survey Special Map 180, with text, 94 p.

COLUMN 12
COVINGTON COUNTY

SYSTEM	SERIES	GROUP, FORMATION, AND MEMBER	LITHO-LOGY	THICKNESS (feet)	THICKNESS (meters)
Tertiary	Oligocene	Bumpnose Limestone		16	5
		Crystal River Formation		121	37
		Yazoo Clay		59	18
		Moodys Branch Formation		59	18
		Lisbon Formation		170	52
		Tallahatta Formation		170	52
		Hatchetigbee Formation		56	17
		Bashi Marl Member		36	11
		Tusahoma Sand		266	81
		Nanafalia Formation		354	108
Cretaceous	Paleocene	Midway Group		334	102
		Clayton Formation		95	29
	Upper	Selma Group undifferentiated		1433	437
		Eutaw Formation		222	68
		Tuscaloosa Group		1115	339
		"Upper Tuscaloosa"		605	184
		"Marine Tuscaloosa"		128	39
		"Lower Tuscaloosa"		382	116
	Lower	Lower Cretaceous undifferentiated		4116	1255
		Cotton Valley Group		1316	401
Jurassic	Upper	Haynesville Formation		170	52
Triassic		Smackover Formation		98	30
		Norphlet Formation		88	27
	Middle	Werner Formation		10	3
	Upper	Eagle Mills Formation		22P	7P

The Covington County section is based on surface exposures, a description of samples from a water well in the town of Florala and the sample descriptions and electric logs from oil and gas test wells (Alabama State Oil and Gas Board Permit Numbers 183 and 1951). Total depths of these two wells are 4,510 and 10,020 feet, respectively.

COLUMN 13
DALLAS COUNTY

SYSTEM	SERIES	GROUP, FORMATION, AND MEMBER	LITHO-LOGY	THICKNESS (feet)	THICKNESS (meters)
Tertiary	Paleocene	Clayton Formation		30	9
		McBryde Ls. Mbr.		112	34
		Pine Barren Mbr.		10-56	3-17
		Prairie Bluff Chalk		180	55
		Ripley Formation		390	119
		Demopolis Chalk		10-1036	3-316
		Arcola Ls. Mbr.		10	3
		Mooreville Chalk		400	122
		Eutaw Formation		390	119
		Tuscaloosa Group		1325	404
Cretaceous	Upper	"Upper Tuscaloosa"		650	198
		"Marine Tuscaloosa"		140	43
		"Lower Tuscaloosa"		535	163
	Lower	Lower Cretaceous undifferentiated		1182	360
		Cotton Valley Group		465	142
	Upper	Haynesville Formation		70	21
		Smackover Formation		20	6
		Norphlet Formation		300	91
	Middle	Werner Formation		106	32

The Tertiary and uppermost Cretaceous units are from surface exposures. The remainder of the column is from adjacent oil and gas test wells (Alabama State Oil and Gas Board Permit Numbers 849 and 1314). Total depths of these two wells are 5,205 and 4,210 feet, respectively.

SELECTED REFERENCES

Column 14

Causey, L. V., Scott, J. C., and Newton, J. G., 1967, Geologic map of Houston County, Alabama: Alabama Geological Survey Special Map 58.

Drennen, C. W., 1953, Reclassification of the outcropping Tuscaloosa Group in Alabama: American Association Petroleum Geologists Bulletin, v. 37, no. 3, p. 522-538.

Moore, D. B., and Joiner, T. J., 1969, A subsurface study of southeastern Alabama: Alabama Geological Survey Bulletin 88, 33 p.

Rheinhardt, Juergen, Gibson, T. G., and others, 1980, Upper Cretaceous and Lower Tertiary geology of the Chattahoochee River Valley, western Georgia and eastern Alabama: American Geological Institute, Excursions in Southeastern Geology, v. II, p. 385-463.

Toulmin, L. D., 1977, Stratigraphic distribution of Paleocene and Eocene fossils in the eastern Gulf Coast region: Alabama Geological Survey Monograph 13, 602 p.

Column 15

Drennen, C. W., 1953, Reclassification of the outcropping Tuscaloosa Group in Alabama: American Association Petroleum Geologists Bulletin, v. 37, no. 3, p. 522-538.

Monroe, W. H., 1941, Notes on deposits of Selma and Ripley age in Alabama: Alabama Geological Survey Bulletin 48, 150 p.





Moore, D. B., and Joiner, T. J., 1969, A subsurface study of southeastern Alabama: Alabama Geological Survey Special Map 58.

Newton, J. G., 1965, Geologic map of Barbour County, Alabama: Alabama Geological Survey Special Map 33.

Rheinhardt, Juergen, Gibson, T. G., and others, 1980, Upper Cretaceous and Lower Tertiary geology of the Chattahoochee River Valley, western Georgia and eastern Alabama: American Geological Institute, Excursions in Southeastern Geology, v. II, p. 385-463.

Toulmin, L. D., 1977, Stratigraphic distribution of Paleocene and Eocene fossils in the eastern Gulf Coast region: Alabama Geological Survey Monograph 13, 602 p.

COLUMN 14
HOUSTON COUNTY

SYSTEM	SERIES	GROUP, FORMATION, AND MEMBER	LITHO-LOGY	THICKNESS (feet)	THICKNESS (meters)
Tertiary	Miocene-Oligocene	Residium		60	18
		Crystal River Formation		46	14
	Eocene	Moodys Branch Formation		46	14
		Lisbon Formation		60	18
		Tallahatta Formation		370	113
		Hatchetigbee Formation		95	29
		Bashi Marl Member		25	8
	Paleocene	Tusahoma Sand		300	92
		Nanafalia Formation		320	98
		Midway Group		375	114
Cretaceous	Upper	Selma Group correlatives		1005	306
		Eutaw Formation		240	73
		Tuscaloosa Group	"Upper Tuscaloosa"	165	50
			"Marine Tuscaloosa"	195	59
			"Lower Tuscaloosa"	210	64
	Lower	Lower Cretaceous undifferentiated		3495	1066
Triassic		Triassic ? undifferentiated		545	166
Ordovician	Lower	Ordovician undifferentiated		545 P	166 P

The Houston County section is based on an oil and gas test well (Alabama State Oil and Gas Board Permit Number 186). Total depth of the well is 8,100 feet.

COLUMN 15
BARBOUR COUNTY

SYSTEM	SERIES	GROUP, FORMATION, AND MEMBER	LITHO-LOGY	THICKNESS (feet)	THICKNESS (meters)
Tertiary	Paleocene	Clayton Formation		200	61
Cretaceous	Upper	Selma Group	Providence Sand	195	59
			Ripley Formation	275	84
			Cusseta Sand Mbr	95	29
			Blufftown Formation	965	294
		Eutaw Formation		190	58
	Lower	Tuscaloosa Group	"Upper Tuscaloosa"	365	111
			"Marine Tuscaloosa"	80	24
			"Lower Tuscaloosa"	310	94
			Lower Cretaceous undifferentiated	1731	528
		Triassic undifferentiated		1140 P	347 P

The Barbour County section is based on a sample description and electric log from an oil and gas test well (Alabama State Oil and Gas Board Permit Number B-321). Total depth of the well is 5,546 feet.

APPALACHIAN PLATEAUS PROVINCE
AND
INTERIOR LOW PLATEAUS PROVINCE

SELECTED REFERENCES

Column 16

Conant, L. C., and Swanson, V. E., 1961, Chattanooga Shale and related rocks of central Tennessee and nearby areas: U.S. Geological Survey Professional Paper 357, 91 p.

Jewell, J. W., 1969, An oil and gas evaluation of north Alabama: Alabama Geological Survey Bulletin 93, 65 p.

Kidd, J. T., 1975, Pre-Mississippian subsurface stratigraphy of the Warrior Basin in Alabama: Gulf Coast Association Geological Societies Transactions, v. 25, p. 20-39.

Kidd, J. T., and Neathery, T. L., 1976, Correlation between Cambrian rocks of the southern Appalachian geosyncline and Interior Low Plateaus: Geology, v. 4, p. 767-769.

Neathery, T. L., and Drahovzal, J. A., 1985, Lithostratigraphy of Upper Ordovician strata in Alabama: Alabama Geological Survey Circular 124, 55 p.

Smith, W. E., 1979, Pennsylvanian stratigraphy of Alabama, *in* The Mississippian and Pennsylvanian (Carboniferous) Systems in the United States: U.S. Geological Survey Professional Paper 1110 I, p. 123 - 136.

Thomas, W. A., 1972, Mississippian stratigraphy of Alabama: Alabama Geological Survey Monograph 12, 121 p.

Column 17

Conant, L. C., and Swanson, V. E., 1961, Chattanooga Shale and related rocks of central Tennessee and nearby areas: U.S. Geological Survey Professional Paper 357, 91 p.

Kidd, J. T., 1975, Pre-Mississippian subsurface stratigraphy of the Warrior Basin in Alabama: Gulf Coast Association Geological Societies Transactions, v. 25, p. 20-39.

Kidd, J. T., and Neathery, T. L., 1976, Correlation between Cambrian rocks of the southern Appalachian geosyncline and Interior Low Plateaus: Geology, v. 4, p. 767-769.

Neathery, T. L., and Drahovzal, J. A., 1985, Lithostratigraphy of Upper Ordovician strata in Alabama: Alabama Geological Survey Circular 124, 55 p.

Smith, W. E., 1979, Pennsylvanian stratigraphy of Alabama, *in* The Mississippian and Pennsylvanian (Carboniferous) Systems in the United States: U.S. Geological Survey Professional Paper 1110 I, p. 123 - 136.

Thomas, W. A., 1972, Mississippian stratigraphy of Alabama: Alabama Geological Survey Monograph 12, 121 p.

Wilson, R. L., 1971, Geology of Jackson County, Alabama: Alabama Geological Survey open-file report, 68 p.

COLUMN 16
MADISON COUNTY

SYSTEM	SERIES	GROUP, FORMATION AND MEMBER	LITHO-LOGY	THICKNESS (feet)	THICKNESS (meters)	
	Lower	Pottsville Formation		200	61	
Mississippian	Upper	Bangor Limestone		400 - 500	122 - 152	
		Hartselle Sandstone		0 - 26	0 - 8	
		Pride Mountain Formation		10 - 23	3 - 7	
		Monteagle Limestone		200 - 220	61 - 67	
		Tuscumbia Limestone		150	46	
Devonian	Lower	Fort Payne Chert		155 - 185	47 - 56	
		Maury Formation		0 - 3	0 - 1	
Silurian		Chattanooga Shale		0 - 23	0 - 7	
		Wayne Group		0 - 15	0 - 5	
		Brassfield Limestone		0 - 35	0 - 11	
	Upper	Sequatchie Formation		0 - 45	0 - 14	
		Leipers Limestone		20 - 60	6 - 18	
Ordovician		Inman Formation		20 - 43	6 - 13	
		Nashville Group		200 - 262	61 - 80	
		Middle	Stones River Group undifferentiated		680 - 780	207 - 238
	Pond Spring Formation			59 - 151	18 - 46	
	Lower	Knox Group undifferentiated		3200 E (977 P)	976 E (298 P)	
	Cambrian	Upper			3200 E (977 P)	976 E (298 P)
		Middle	Ketona Dolomite		750 E	229 E
			Conasauga Formation		1400 E	427 E
Pre-cambrian		Lower	Rome Formation		330E	101 E
		Granitic rock				

The Pennsylvanian and Mississippian section was derived from published information. The Ordovician to Devonian section was based on published information and sample logs for several wells (Alabama State Oil and Gas Board Permit Numbers 1011, 1563, 1630, and 1640). The Cambrian section is based on data from wells drilled in Cullman and Morgan Counties, Alabama, and on published data.

COLUMN 17
JACKSON COUNTY

SYSTEM	SERIES	GROUP, FORMATION AND MEMBER	LITHO-LOGY	THICKNESS (feet)	THICKNESS (meters)						
Pennsyl- vanian	Lower	Pottsville Formation		100 - 394	30 - 120						
Mississippian	Upper	Pennington Formation		228 - 445	70 - 136						
		Bangor Limestone		200 - 400	61 - 122						
		Monteagle Limestone		200 - 300	61 - 91						
		Tuscumbia Limestone		200	61						
		Fort Payne Chert		150	46						
	Lower	Chattanooga Shale		12 - 40	4 - 12						
Devonian	Upper	Red Mountain Formation equivalent		95 - 180	29 - 55						
		Sequatchie Formation		66	20						
		Leipers Limestone		56	17						
		Inman Formation		320	98						
		Nashville Group		320	98						
	Middle	Stones River Group		675	206						
			Pond Spring Formation		239	73					
		Lower	Knox Group undifferentiated		3200 E	976 E					
	Cambrian	Upper		Ketona Dolomite		394 E	120 E				
								Middle	Conasauga Formation		2000 E
Lower		Rome Formation			590 E	180 E					
Pre-cam- brian				Granitic rock							

The Devonian-Pennsylvanian section was derived from published information and from the Geological Survey of Alabama open-file geologic map of Jackson County. The Ordovician-Devonian section was based on published information and the sample log for an oil and gas test well (Alabama State Oil and Gas Board Permit Number 2059). The Cambrian-Precambrian section was based on data from wells drilled in Cullman and Morgan Counties, Alabama, and on published data.

SELECTED REFERENCES

Column 18

Burchard, E. F., and Andrews, T. G., 1947, Iron ore outcrops of the Red Mountain Formation in northeast Alabama: Alabama Geological Survey Special Report 19, 375 p.

Conant, L. C., and Swanson, V. E., 1961, Chattanooga Shale and related rocks of central Tennessee and nearby areas: U.S. Geological Survey Professional Paper 357, 91 p.

Kidd, J. T., 1975, Pre-Mississippian subsurface stratigraphy of the Warrior Basin in Alabama: Gulf Coast Association Geological Societies Transactions, v. 25, p. 20-39.

Neathery, T. L., Smith, W. E., Clarke, O. M., Jr., and Szabo, M. W., 1968, Geology and mineral resources of Etowah County, Alabama: Alabama Geological Survey open-file report, 281 p.

Neathery, T. L., and Drahovzal, J. A., 1985, Lithostratigraphy of Upper Ordovician strata in Alabama: Alabama Geological Survey Circular 124, 55 p.

Smith, W. E., 1979, Pennsylvanian stratigraphy of Alabama, *in* The Mississippian and Pennsylvanian (Carboniferous) Systems in the United States: U.S. Geological Survey Professional Paper 1110 I, p. 123-136.

Thomas, W. A., 1972, Mississippian stratigraphy of Alabama: Alabama Geological Survey Monograph 12, 121 p.

Column 19

Burchard, E. F., and Andrews, T. G., 1947, Iron ore outcrops of the Red Mountain Formation in northeast Alabama: Alabama Geological Survey Special Report 19, 375 p.

Butts, Charles, 1926, The Paleozoic rocks, *in* Geology of Alabama: Alabama Geological Survey Special Report 14, p. 41-230.

Conant, L. C., and Swanson, V. E., 1961, Chattanooga Shale and related rocks of central Tennessee and nearby areas: U.S. Geological Survey Professional Paper 357, 91 p.

Faust, R. J., 1984, Geology of Blount County, Alabama: Alabama Geological Survey Special Map 159, 29 p.

Kidd, J. T., 1975, Pre-Mississippian subsurface stratigraphy of the Warrior Basin in Alabama: Gulf Coast Association Geological Societies Transactions, v. 25, p. 20-39.

Kidd, J. T., and Neathery, T. L., 1976, Correlation between Cambrian rocks of the southern Appalachian geosyncline and Interior Low Plateaus: Geology, v. 4, p. 767-769.

Neathery, T. L., and Drahovzal, J. A., 1985, Lithostratigraphy of Upper Ordovician strata in Alabama: Alabama Geological Survey Circular 124, 55 p.

Smith, W. E., 1979, Pennsylvanian stratigraphy of Alabama, *in* The Mississippian and Pennsylvanian (Carboniferous) Systems in the United States: U.S. Geological Survey Professional Paper 1110 I, p. 123-136.

Thomas, W. A., 1972, Mississippian stratigraphy of Alabama: Alabama Geological Survey Monograph 12, 121 p.

COLUMN 18
DE KALB COUNTY AND
WESTERN ETOWAH COUNTY

SYSTEM	SERIES	GROUP, FORMATION, AND MEMBER	LITHO- LOGY	THICKNESS (feet)	THICKNESS (meters)																																																																	
Pennsylvanian	Lower	Pottsville Formation		25 - 1400	8 - 427																																																																	
						Mississippian	Upper	Parkwood and Pennington Formations undifferentiated		150 - 550	46 - 168	Bangor Limestone		295 - 550	90 - 168	Hartselle Sandstone	Monteagle Limestone	0 - 75	0 - 23	Pride Mountain Formation	0 - 75	0 - 23	Tuscumbia Limestone		15 - 25	4 - 8	Lower	Fort Payne Chert	100 - 200	30 - 61	Maury Formation	3 - 6	1 - 2	Chattanooga Shale	3 - 50	1 - 15					Devonian						Silurian	Upper	Red Mountain Formation		450 - 700	137 - 213	Lower	Ordovician	Upper	Sequatchie Formation		200	61	Middle	Nashville and Stones River Groups undifferentiated or Chickamauga Limestone		200 - 1100	61 - 335	Lower	Knox Group undifferentiated		3500	1067	Cambrian
Mississippian	Upper	Parkwood and Pennington Formations undifferentiated		150 - 550	46 - 168																																																																	
		Bangor Limestone		295 - 550	90 - 168																																																																	
		Hartselle Sandstone	Monteagle Limestone	0 - 75	0 - 23																																																																	
		Pride Mountain Formation		0 - 75	0 - 23																																																																	
		Tuscumbia Limestone		15 - 25	4 - 8																																																																	
		Lower	Fort Payne Chert	100 - 200	30 - 61																																																																	
	Maury Formation		3 - 6	1 - 2																																																																		
	Chattanooga Shale		3 - 50	1 - 15																																																																		
	Devonian																																																																					
Silurian	Upper	Red Mountain Formation		450 - 700	137 - 213																																																																	
	Lower																																																																					
Ordovician	Upper	Sequatchie Formation		200	61																																																																	
	Middle	Nashville and Stones River Groups undifferentiated or Chickamauga Limestone		200 - 1100	61 - 335																																																																	
	Lower	Knox Group undifferentiated		3500	1067																																																																	
Cambrian	Upper																																																																					

The entire section was based on data in published literature and cited references.

COLUMN 19
BLOUNT COUNTY

SYSTEM	SERIES	GROUP, FORMATION, AND MEMBER	LITHO-LOGY	THICKNESS (feet)	THICKNESS (meters)
Pennsylvanian	Lower	Pottsville Formation		33 - 984	10 - 300
Mississippian	Upper	Parkwood Formation		0 - 400	0 - 122
		Bangor Limestone		400 - 700	122 - 213
		Hartselle Sandstone		24 - 100	7 - 30
		Pride Mountain Formation		50 - 125	
		Tuscumbia Limestone		200 - 250	61 - 76
		Fort Payne Chert		150 - 210	46 - 64
	Lower	Chattanooga Shale		1 - 45	1 - 14
		Red Mountain Formation		158 - 350	48 - 107
	Upper	Richmond, Maysville and Eden Groups undifferentiated		90	27
		Nashville Group		66 - 110	20 - 34
Ordovician	Middle	Stones River Group		820	250
	Lower	Knox Group undifferentiated		3070	936
	Upper				
		Ketona Dolomite		215	66
Cambrian	Middle	Conasauga Formation		2015	614
	Lower	Rome Formation		410	125
		Granitic rock		50P	15P

The Pennsylvanian section was based on published literature and cited references. The pre-Pennsylvanian to Middle Ordovician section was based on cited references and the sample log for an oil and gas test well (Alabama State Oil and Gas Board Permit Number 2795). The section of Middle Ordovician and older rocks was based on the sample log of Permit Number 2795. Total depth of the well is 8,430 feet.

SELECTED REFERENCES

Column 20	Column 21
Conant, L. C., and Swanson, V. E., 1961, Chattanooga Shale and related rocks of central Tennessee and nearby areas: U.S. Geological Survey Professional Paper 357, 91 p.	Butts, Charles, 1926, The Paleozoic rocks, <i>in</i> Geology of Alabama: Alabama Geological Survey Special Report 14, p. 41-230.
Faust, R. J., and Jefferson, P. D., 1980, Geology and water availability of Cullman County, Alabama: Alabama Geological Survey Special Map 145.	Conant, L. C., and Swanson, V. E., 1961, Chattanooga Shale and related rocks of central Tennessee and nearby areas: U.S. Geological Survey Professional Paper 357, 91 p.
Jewell, J. W., 1969, An oil and gas evaluation of north Alabama: Alabama Geological Survey Bulletin 93, 65 p.	Dodson, C. L., and Harris, W. F., Jr., 1965, Geology and ground-water resources of Morgan County, Alabama: Alabama Geological Survey Bulletin 76, 90 p.
Kidd, J. T., 1975, Pre-Mississippian subsurface stratigraphy of the Warrior Basin in Alabama: Gulf Coast Association Geological Societies Transactions, v. 25, p. 20-39.	Jewell, J. W., 1969, An oil and gas evaluation of north Alabama: Alabama Geological Survey Bulletin 93, 65 p.
Lundin, R. F., and Newton, G. D., 1970, Ostracoda and the Silurian stratigraphy of northwestern Alabama: Alabama Geological Survey Bulletin 95, 64 p.	Kidd, J. T., 1975, Pre-Mississippian subsurface stratigraphy of the Warrior Basin in Alabama: Gulf Coast Association Geological Societies Transactions, v. 25, p. 20-39.
Neathery, T. L., and Drahovzal, J. A., 1985, Lithostratigraphy of Upper Ordovician strata in Alabama: Alabama Geological Survey Circular 124, 55 p.	Kidd, J. T., and Neathery, T. L., 1976, Correlation between Cambrian rocks of the southern Appalachian geosyncline and Interior Low Plateaus: Geology, v. 4, p. 767-769.
Smith, W. E., 1979, Pennsylvanian stratigraphy of Alabama, <i>in</i> The Mississippian and Pennsylvanian (Carboniferous) Systems in the United States: U.S. Geological Survey Professional Paper 1110 I, p. 123-136.	Lundin, R. F., and Newton, G. D., 1970, Ostracoda and the Silurian stratigraphy of northwestern Alabama: Alabama Geological Survey Bulletin 95, 64 p.
Thomas, W. A., 1972, Mississippian stratigraphy of Alabama: Alabama Geological Survey Monograph 12, 121 p.	Neathery, T. L., and Drahovzal, J. A., 1985, Lithostratigraphy of Upper Ordovician strata in Alabama: Alabama Geological Survey Circular 124, 55 p.
	Thomas, W. A., 1972, Mississippian stratigraphy of Alabama: Alabama Geological Survey Monograph 12, 121 p.

COLUMN 20
CULLMAN COUNTY

SYSTEM	SERIES	GROUP, FORMATION, AND MEMBER	LITHO-LOGY	THICKNESS (feet)	THICKNESS (meters)
Pennsylvanian	Lower	Pottsville Formation		360	110
Mississippian	Upper	Bangor Limestone		520	158
		Hartselle Sandstone		15	4
		Pride Mountain Fm.-Monteagle Ls. undiff.		85	26
		Tuscumbia Limestone		260	79
	Lower	Fort Payne Chert		160	49
Silurian		Chattanooga Shale		24	7
		Silurian undifferentiated		22	7
	Upper	Richmond Group ?		50	15
		Elkmont Limestone		130	40
		Hermitage Formation		50	15
Ordovician	Middle	Stones River Group		695	212
		Pond Spring Formation		150	46
	Lower	Knox Group undifferentiated		3215	980
Cambrian	Upper	Ketona Dolomite		760	232
	Middle	Conasauga Formation		1400	427
Pre-cambrian	Lower	Rome Formation		290	88
		Basement (granitic gneiss)		80P	24P

The entire column was based on the electric log and sample description of an oil and gas test well (Alabama State Oil and Gas Board Permit Number 1689). Total depth of the well is 8,270 feet.

COLUMN 21
MORGAN COUNTY

SYSTEM	SERIES	GROUP, FORMATION, AND MEMBER	LITHO-LOGY	THICKNESS (feet)	THICKNESS (meters)
Mississippian	Upper	Bangor Limestone		180	55
		Hartselle Sandstone		4	1
		Pride Mountain Fm. Monteagle Ls. undiff.		170	52
		Tuscumbia Limestone		218	66
Devonian	Lower	Fort Payne Chert		190	58
		Chattanooga Shale		18	5
		Silurian undifferentiated		100	30
Ordovician	Middle	Ordovician undifferentiated		226	69
		Stones River Group		774	236
	Lower	Knox Group undifferentiated		3500	1067
Cambrian	Upper				
	Ketona Dolomite		690	210	
	Middle	Conasauga Formation		1250	381
Precambrian	Lower	Rome Formation		320	98
		Basement (granite)		28P	8P

The entire column was based on the electric log and sample description of an oil and gas test well (Alabama State Oil and Gas Board Permit Number 2794). Total depth of the well is 7,668 feet.

VALLEY AND RIDGE PROVINCE

SELECTED REFERENCES

Column 22

Butts, Charles, 1926, The Paleozoic rocks, *in* Geology of Alabama: Alabama Geological Survey Special Report 14, p. 41-230.

Kidd, J. T., 1979, Areal geology of Jefferson County, Alabama: Alabama Geological Survey Atlas 15, 89 p.

Tucker, W. E., and Kidd, R. E., 1973, Appendix G, Sample log U.S. Steel Corporation Disposal Well No. 1, Fairfield, Alabama, *in* Deep-well disposal in Alabama: Alabama Geological Survey Bulletin 104, p. 191-208.

Column 23

Causey, L. V., 1965a, Geologic map of Cherokee County, Alabama: Alabama Geological Survey Special Map 39.

———, 1965b, Geology and ground-water resources of Cherokee County, Alabama, a reconnaissance: Alabama Geological Survey Bulletin 79, 63 p.

Neathery, T. L., Smith, W. E., Clarke, O. M., Jr., and Szabo, M. W., 1968, Geology and mineral resources of Etowah County, Alabama: Alabama Geological Survey open-file report, 281 p.

Thomas, W. A., 1972, Mississippian stratigraphy of Alabama: Alabama Geological Survey Monograph 12, 121 p.

SYSTEM	SERIES	GROUP, FORMATION, AND MEMBER		LITHO-LOGY	THICKNESS (feet)	THICKNESS (meters)
Pennsylvanian	Lower	Pottsville Formation			2788 - 9496	850 - 2895
Mississippian	Upper	Parkwood Formation			900 - 2500	274 - 762
		Bangor Limestone	Floyd Shale			
		Hartselle Sandstone				
		Pride Mountain Formation				
		Tuscumbia Limestone				
	Lower	Fort Payne Chert			88 - 200	27 - 61
		Maury Formation				
		Chattanooga Shale				
		Frog Mountain Sandstone				
Devonian						
Silurian		Red Mountain Formation		200 - 500	61 - 152	
Ordovician	Upper	Sequatchie Formation			0 - 3	0 - 1
	Middle	Chickamauga Limestone			260 - 815	79 - 248
		Attalla Chert Conglomerate Mbr.			0 - 50	0 - 15
	Lower	Knox Group undifferentiated			1475 - 2950	450 - 899
Cambrian	Upper	Ketona Dolomite			0 - 600	0 - 180
	Middle	Conasauga Formation			1100 - 1900	335 - 579

This column is based on exposures in the Valley and Ridge part of Jefferson County northwest of the Helena fault and the sample log for a disposal well drilled by U.S. Steel in sec. 10, T. 18 S., R. 4 W. Total depth of the disposal well is 6,060 feet.

COLUMN 22
JEFFERSON COUNTY

COLUMN 23
NORTHEASTERN ETOWAH AND
NORTHWESTERN CHEROKEE COUNTIES

	SERIES	GROUP, FORMATION, AND MEMBER	LITHO-LOGY	THICKNESS (feet)	THICKNESS (meters)
Pennsylvanian	Lower	Pottsville Formation		23 - 200	7 - 61
Mississippian	Upper	Parkwood and Pennington Formations undifferentiated		500	152
		Bangor Limestone and Monteagle Limestone undifferentiated		880	268
		Tuscumbia Limestone		100	30
		Fort Payne Chert		150	46
		Maury Formation		3 - 6	1 - 2
	Lower	Chattanooga Shale		20 - 40	6 - 12
		Red Mountain Formation		395 - 590	120 - 180
	Upper	Sequatchie Formation		100 - 200	30 - 60
	Middle	Chickamauga Limestone		700 - 1000	213 - 305
		Attalla Chert Conglomerate Mbr.		0 - 108	0 - 33
Ordovician	Lower	Knox Group undifferentiated		3500	1067
	Upper				
	Middle	Conasauga Formation		1400 +	427 +

The entire column is based on data in published literature and cited references.

SELECTED REFERENCES

Column 24

Butts, Charles, 1926, The Paleozoic rocks, *in* Geology of Alabama: Alabama Geological Survey Special Report 14, p. 41-230.

Causey, L. V., 1965a, Geologic map of Cherokee County, Alabama: Alabama Geological Survey Special Map 39.

———, 1965b, Geology and ground-water resources of Cherokee County, Alabama, a reconnaissance: Alabama Geological Survey Bulletin 79, 63 p.

Drahovzal, J. A., and Neathery, T. L., 1971, Middle and Upper Ordovician stratigraphy of the Alabama Appalachians, *in* The Middle and Upper Ordovician of the Alabama Appalachians: Alabama Geological Society Guidebook, Ninth Annual Field Trip, p. 1-62.

Drahovzal, J. A., and Thomas, W. A., 1977, Pre-Mississippian sandstones in the interior structures of the Appalachian fold and thrust belt of Eastern Alabama, *in* Bearce, D. N., ed., Cambrian and Devonian stratigraphic problems of eastern Alabama: Alabama Geological Society Guidebook, 15th Annual Field Conference, p. 29-36.

Mack, G. H., 1980, Stratigraphy and depositional environments of the Chilhowee Group (Cambrian) in Georgia and Alabama: American Journal of Science, v. 280, p. 497-517.

Neathery, T. L., Smith, W. E., Clarke, O. M., Jr., and Szabo, M. W., 1968, Geology and mineral resources of Etowah County, Alabama: Alabama Geological Survey open-file report, 281 p.

Thomas, W. A., 1972, Mississippian stratigraphy of Alabama: Alabama Geological Survey Monograph 12, 121 p.

Thomas, W. A., and Drahovzal, J. A., eds., 1974, The Coosa deformed belt in the Alabama Appalachians: Alabama Geological Society Guidebook, 12th Annual Field Trip, 98 p.

Warman, J. C., and Causey, L. V., 1962, Geologic map of Calhoun County, Alabama: Alabama Geological Survey Special Map 17.

SYSTEM	SERIES	GROUP, FORMATION, AND MEMBER		LITHO-LOGY	THICKNESS (feet)	THICKNESS (meters)
Pennsylvanian	Lower	Pottsville Formation			300 ±	91 ±
		Parkwood Formation			1269	387
Mississippian	Upper	Floyd Shale			1200 - 2000	366 - 610
		Fort Payne Chert			15 - 175	4 - 53
Devonian	Lower	Maury Formation			0 - 3	0 - 1
		Frog Mountain Sandstone			0 - 200	0 - 61
	Upper	Sequatchie Formation			0 - 120	0 - 37
		Athens Shale			0 - 21	0 - 6
	Middle	Colvin Mountain Sandstone			0 - 200	0 - 61
		Greensport Formation			230 - 330	70 - 100
	Lower	Little Oak Limestone			0 - 33	0 - 10
		Lenoir Limestone			0 - 250	0 - 76
	Lower	Newala and Longview Limestones			660	201
		Chepultepec Dolomite			1245	380
	Upper	Copper Ridge Dolomite			1245	380
		Conasauga Formation			1900	579
	Lower	Rome Formation			700 - 1000	213 - 305
		Shady Dolomite			370 - 600	113 - 183
	Lower	Weisner Formation			492	150
		Wilson Ridge Formation			603	184
	Lower	Nichols Formation			410	125
		Cochran Formation			233 +	71 +

COLUMN 24 CALHOUN, SOUTHEASTERN CHEROKEE AND SOUTHEASTERN ETOWAH COUNTIES

The Pennsylvanian part of the column is based on the westernmost part of Calhoun County. The Mississippian part is based on cited references on all three counties. The Devonian part is based on cited references and is present in all three counties. The Cambrian-Ordovician section is based on exposures in all three counties. The best exposures of the Ordovician are on Colvin and Greens Creek Mountains in Calhoun and Etowah Counties.

SELECTED REFERENCES

Column 25



Butts, Charles, 1926, The Paleozoic rocks, *in* Geology of Alabama: Alabama Geological Survey Special Report 14, p. 41-230.

Raymond, D. E., 1985, New information on lower Paleozoic stratigraphy of the Alabama fold and thrust belt: Geological Society of America Southeastern Section, 34th Annual Meeting, Abstracts with Programs, v. 17, no. 2, p. 130.

COLUMN 25

SHELBY COUNTY
SUBSURFACE

SYSTEM	SERIES	GROUP, FORMATION, AND MEMBER		LITHO- LOGY	THICKNESS (feet)	THICKNESS (meters)
Mississippian		Floyd Shale			779	238
		Tuscumbia Limestone- Fort Payne Chert undiff.			64	20
		Fort Payne Chert			90	27
Ordovician	Middle	Athens Shale			29	9
		Little Oak-Lenoir Limestones undifferentiated			237	72
	Lower	Knox Group	Newala Limestone		531	162
			Longview Limestone		407	124
			Chepultepec Dolomite		778	237
						2801
	Cambrian	Upper	Copper Ridge Dolomite			1085
Ketona Dolomite equivalent				834	254	
Ordovician		Middle	Lenoir Limestone			362
	Lower	Knox Group	Newala Limestone		1654	504
			Longview Limestone		450	137
			Chepultepec Dolomite		966	294
					4290	1307
Cambrian	Upper	Copper Ridge Dolomite			1220	372

SYSTEM	SERIES	GROUP, FORMATION, AND MEMBER	LITHO- LOGY	THICKNESS (feet)	THICKNESS (meters)
Cambrian	Upper	Ketona Dolomite		929	283
	Middle	Conasauga Formation		271	83
		Rome Formation		2953	900
	Lower	Shady Dolomite		338	103
	Middle	Conasauga Formation		3028	923

This column for Shelby County is based on the density log and sample log for an oil and gas test well (Alabama State Oil and Gas Board Permit Number 3518). Total depth of the well is 17,005 feet. The well is in the vicinity of the Page Springs anticline in sec. 15, T. 20 S., R. 1 E.

SELECTED REFERENCES

Column 26

Butts, Charles, 1926, The Paleozoic rocks, *in* Geology of Alabama: Alabama Geological Survey Special Report 14, p. 41-230.

——— 1927, Description of Bessemer-Vandiver quadrangles [Alabama]: U.S. Geological Survey Geologic Atlas, Folio 221, 22 p.

——— 1940, Description of Montevallo-Columbiana quadrangles [Alabama]: U.S. Geological Survey Geologic Atlas, Folio 226, 20 p.

Drahovzal, J. A., and Neathery, T. L., 1971, Middle and Upper Ordovician stratigraphy of the Alabama Appalachians, *in* The Middle and Upper Ordovician of the Alabama Appalachians: Alabama Geological Society Guidebook, Ninth Annual Field Trip, p. 1-62.

Raymond, D. E., 1985, New information on lower Paleozoic stratigraphy of the Alabama fold and thrust belt: Geological Society of America Southeastern Section, 34th Annual Meeting, Abstracts with Programs, v. 17, no. 2, p. 130.

Thomas, W. A., 1972, Mississippian stratigraphy of Alabama: Alabama Geological Survey Monograph 12, 121 p.

COLUMN 26
BIBB AND SHELBY COUNTIES

SYSTEM	SERIES	GROUP, FORMATION, AND MEMBER	LITHO-LOGY	THICKNESS (feet)	THICKNESS (meters)
Pennsylvanian	Lower	Pottsville Formation		4920 - 9000 ±	1500 - 2745 ±
Mississippian	Upper	Parkwood Formation		470 - 2000 +	143 - 610 +
		Floyd Shale		800 - 2000	244 - 610
Devonian	Lower	Fort Payne Chert		0 - 200	0 - 61
		Chattanooga Shale		0 - 7	0 - 2
Ordovician	Middle	Frog Mountain Sandstone		0 - 75	0 - 23
		Little Oak Limestone		100 - 15	30 - 5
		Lenoir Limestone		400 - 700	122 - 213

SYSTEM	SERIES	GROUP, FORMATION, AND MEMBER	LITHO-LOGY	THICKNESS (feet)	THICKNESS (meters)
Ordovician	Lower	Newala and Longview Limestones		1200 - 1400	366 - 427
		Chepultepec Dolomite		1100	335
Cambrian	Upper	Copper Ridge Dolomite		1750	534
		Bibb Dolomite		275 - 500	84 - 152
		Ketona Dolomite		275 - 1000	84 - 305
		Brierfield Dolomite		0 - 1250	0 - 381
	Middle	Conasauga Formation		0 - 1500	0 - 457
		Rome Formation		1000 Exp	305 Exp

This is a composite column based on exposures throughout the eastern Valley and Ridge in both counties.

Published by
GEOLOGICAL SURVEY OF ALABAMA
Serving Alabama since 1848
Ernest A. Mancini, State Geologist

For additional copies write:
Publication Sales Office
P. O. Box 1424
Tuscaloosa, Alabama 35486-1424
Phone 205/349-2852